

Research Article

Determination of toxigenic potential of *Fusarium* species occurring on sorghum and maize grains produced in Karnataka, India by using Thin Layer Chromatography

M Y Sreenivasa ¹, B T Diwakar ², Adkar Purushothama Charith Raj ³, Regina Sharmila Dass ³, K A Naidu ² and G R Janardhana ^{3,*}

¹ Department of studies in Microbiology, University of Mysore, Manasagangotri, Mysore- 570 006, Karnataka, India; ² Department of Biochemistry and Nutrition, Central Food Technological Research Institute, Mysore- 570 020, Karnataka, India; ³ Mycology and Phytopathology Laboratory, Department of studies in Botany and Microbiology, University of Mysore, Manasagangotri, Mysore- 570 006, Karnataka, India..

Corresponding author

Email:

grjbelur@gmail.com

Phone No:

091-821-2419763

Fax No:

091-821-2419759

Editor

Ganesh Kumar Agrawal

DOI

dx.doi.org/10.3126/ijls.v6

i1.5883

ABSTRACT

In the present study, the fumonisin producing ability of *Fusarium* species from different geographic regions of Karnataka was investigated. Sorghum and maize samples were examined for the presence of *Fusarium* species. Mycological studies confirmed the occurrence of *Fusarium* species such as *Fusarium verticillioides* (18), *F. proliferatum* (2) and *F. anthophilum* (2) isolates. Efficacy to produce fumonisins on maize patties was carried out by using thin-layer chromatography. Fumonisin B₁ and B₂ were detected as a red-purple spots with an R_f value of 6.1 and 6.3 respectively. The TLC analysis showed that 17 of 18 *F. verticillioides* isolates, 02 of 02 *F. proliferatum* isolates and 01 of 02 *F. anthophilum* isolates produced fumonisins. On the chromatograms, identical colored spots for standards and samples with same R_f value were detected. Early detection of toxigenic *Fusarium* species is important to prevent the entry of these toxic substances into the food chains.

Key words: Fumonisin; *Fusarium sp.*; Maize; mycotoxin; Sorghum

INTRODUCTION

The FAO estimated that each year, 25% to 50% of the world's food crops are contaminated by mycotoxins (Fandohan *et al.* 2003). *Fusarium* species are known to produce a chemically diverse array of mycotoxins such as diacetoxyscirpenol, deoxynivalenol, nivalenol, T-2 toxin, zearalenone, fumonisins, fusarin C, beauvericin, moniliformin, and fusaproliferin (Glenn, 2007; Logrieco *et al.* 2002). Fumonisin is a family of mycotoxins produced primarily by *Fusarium verticillioides* (Saccardo) Nirenberg (= *F. moniliformae*), *F. proliferatum* (Matsushima) Nirenberg, *F. anthophilum* (A. Braun) Wollenweber and others. Although 28 structurally related fumonisin analogues have been identified, only fumonisin B₁ (FB₁) and B₂ (FB₂) occur in abundant levels. Generally FB₁ represents 70-80% of the total fumonisin levels and FB₂ accounts for 15-25% when cultured on maize, rice or in liquid

medium (Rheeder *et al.* 2002). *F. verticillioides* and *F. proliferatum* are the main source of fumonisins, a health risk mycotoxin, contaminating the agro-products (Morales-Rodriguez *et al.* 2007).

Contamination of food commodities by fumonisins has become a serious food safety problem throughout the world (Munkovold and Desjardins 1997). Fumonisin has emerged as a highly visible animal and human health safety concern since they have been associated with many animal diseases (Gelderblom *et al.* 2001). Consumption of corn highly contaminated with fumonisins has been associated with increased risk of human oesophageal cancer in some regions of South Africa (Rheeder *et al.* 1992) and China (Chu and Li 1994). In India, field surveys revealed the occurrence of FB₁ in many samples collected from markets and households in maize and sorghum-growing regions of Karnataka (Sreenivasa *et al.* 2008a; Nayaka *et al.* 2010), Andhra Pradesh (Bhat *et al.* 1997) and Harayana (Jindal *et al.* 1999).

Alarming high levels of fumonisins have also been detected in maize and sorghum (Chhatterjee and Mukherjee, 1994; Jindal et al. 1999). An epidemiological survey conducted in Karnataka and Andhra Pradesh during 1997 revealed that consumption of mouldy grains affected 1424 people from 27 villages (Shetty and Bhat, 1997).

Some of the existing chromatographic methods for detecting fumonisin includes thin-layer chromatography, liquid chromatography, gas chromatography-mass spectroscopy and high-performance liquid chromatography with an electrospray mass spectrometer. Among the available methods, most commonly used method is thin layer chromatographic technique (TLC) as initial screening method. The main objectives of the present investigation were to screen the cultures of *F. verticillioides*, *F. proliferatum* and *F. anthophilum* species isolated from maize and sorghum for their ability to produce fumonisin B₁ and B₂.

MATERIALS AND METHODS

Fusarium strains for fumonisin production

F. verticillioides (18), *F. proliferatum* (2) and *F. anthophilum* (2) isolates were originally isolated from maize and sorghum samples. These isolates were identified up to the species level based on the micro-morphological features using fungal keys and manuals (Leslie and Summerell 2006) and confirmed with molecular markers (Sreenivasa et al. 2006, Sreenivasa et al. 2008b). Further, all the species were evaluated for their ability to produce fumonisins B₁ and B₂ in maize patty cultures.

Preparation of maize patty culture

Preparation of maize patty cultures, inoculation and incubation was performed as per the protocol described by Vismer et al. (2004). Whole maize kernels were ground in a fire proof mixer grinder to a fine powder and 30g was taken in Pyrex Petri dishes (100mm x 18mm) and 30ml of distilled water was added. The preparation was autoclaved at 121 °C for 1 h and allowed to stand overnight. The sterilization was repeated next day for 1 h (Figure-1A). All *Fusarium* species were inoculated on PDA and incubated at 26±2 °C for 4 days. One ml (1x10⁶conidia/ml) of conidial suspension of each fungal isolate was prepared in sterile distilled water and was used to inoculate maize patty in a culture plates. Patties were incubated in dark at 28 ± 2 °C for 4 weeks (Figure-1B), after which they were dried in

an hot air oven at 60 °C for 8-10hrs. Harvested dry patties were ground in a mixer grinder (Kenstar, Classique, MG-9605A) to a fine meal and used for fumonisin analysis.

Clean up with Sep-Pak C₁₈ column

10g of maize patty culture sample was weighed and transferred into a 250ml beaker and mixed with 50ml of acetonitrile/water (50/50, v/v). The beaker was covered with aluminum foil and shaken for 30min. The mixture was filtered through Whatman No. 4 filter paper. The Sep-Pak C₁₈ column was connected to peristaltic pump and was preconditioned with by rinsing with 2ml ACN followed by 1% KCl. Two ml of filtered extract was added to 6ml of 1% KCl and loaded into a C₁₈ clean-up column. The solution was then forced through the Sep-Pak C₁₈ column at a flow rate of 1ml/min. The column was rinsed with 2ml of 1% KCl followed by 2ml of ACN/ H₂O (15+85, v/v). The rinses were discarded, and air was forced through the column to expel all the rinse solution. Finally FB₁ and FB₂ were eluted from the column with 2ml of ACN/ H₂O (70+30, v/v). These elutants were evaporated to complete dryness under a gentle stream of nitrogen and dissolved in 100µl of ACN/ H₂O (50+50, v/v) for TLC analysis.

Thin layer chromatographic analysis

TLC was performed as per the protocol described by (Bially et al. 2005) with some modifications. The samples (5-10µl) was spotted on a thin-layer chromatographic (TLC) plates (20 x 20cm, normal phase) (ALUGRAM® SIL G/UV₂₅₄, Macherey-Nagel, Germany) pre-heated at 110 °C for 10min. The FB₁ and FB₂ (100ng/ml) standards from the stock solutions were spotted on TLC plate. Separation was carried out in 1-butanol-acetic acid-water (20+10+10, v/v/v) as mobile phase. After separation, the plates were dried at room temperature, sprayed with p-anisaldehyde (0.5% in MeOH/sulfuric acid/acetic acid, 90/5/1, v/v/v) and heated at 100 °C for 5min. The R_f value is the ratio the spot moved compared to the distance the solvent moved from the starting point. The detection limit of fumonisins in samples on the TLC plate was 0.1µg/ml.

RESULTS AND DISCUSSION

Fusarium species are among the most studied phytopathogenic fungi, with several species causing diseases on maize, wheat, barley, and other food and feed grains (Glenn, 2007). Wheat, maize and other grain crop residues are generally regarded as the




Figure 1. Maize patty cultures. A: Pyrex petri dishes with maize patty (before inoculation), B: Fusarial growth on maize patties after 4 weeks of incubation.

primary source of inoculum for *Fusarium* species (Shaner, 2003). Decreased yield, as well as diminished quality and value of the grain, results in significant worldwide economic losses. Although fumonisins are a relatively novel class of *Fusarium* toxins, they have attracted both social experts and scientist's attention because of their high health risk potentials (Nelson *et al.* 1992). In some regions of Africa, Asia and other parts of the world, high levels of fumonisins have been constantly detected and recorded on maize (Shephard *et al.* 1996). Many isolates of *F. verticillioides* and *F. proliferatum* isolated from maize and sorghum and other substrates from different geographic locations of North America, Africa, Asia and Australia (Ghianian *et al.* 2005). As identification of *Fusarium* species is critical to predict the potential mycotoxigenic risk of the isolates, there is a need for accurate and complementary tools which permit a

rapid, sensitive and reliable specific diagnosis of potential fumonisin producing *Fusarium* species.

The TLC analysis showed that 17 of 18 *F. verticillioides* isolates, 02 of 02 *F. proliferatum* isolates and 01 of 02 *F. anthophilum* isolates produced fumonisins. Fumonisin B₁ and B₂ were detected as a red-purple spots with an R_f value of 6.1 and 6.3 respectively (Figure 2). On the chromatograms, identical colored spots for standards and samples with same R_f value were detected. The TLC method failed to measure toxin concentrations as accurately as HPLC because the TLC method estimates sample concentration by comparison to standards. The concentration determined by TLC can only be as precise as the number of standards used, and therefore the measured concentration is dependent upon the concentration difference between standards. In view of this, the same samples were confirmed for the presence and quantification of FB1 and FB2 with




Figure 2. Thin-layer chromatographic plate showing the red-purple spots of fumonisins; Lane 1: Standard fumonisin B₁ (Rf - 6.1); Lane 2: Standard fumonisin B₂ (Rf - 6.3); Lane 3 – 8: Positive maize patty samples inoculated with testing *Fusarium* species for Fumonisin; Lane 9: Negative control.

HPLC technique (Sreenivasa *et al.* unpublished data). The two species of *Fusarium* viz., *F. verticillioides* and *F. proliferatum* are the most prolific fumonisin producers and are known to produce fumonisins from 17,900µg/g to 31,000µg/g of FB₁ (Rheeder *et al.* 2002). The production of fumonisins by different strains of *F. proliferatum* varies widely. Some studies have shown that only low or even no fumonisin production (Tseng *et al.* 1995; Da silva *et al.* 2004), while other study has shown that, *F. verticillioides* can produce high levels of fumonisins (Nelson *et al.* 1992). The natural occurrence of fumonisin B₁ in Indian sorghum, where the contamination levels ranging from 0.01-5.0 mg/kg and 0.15-0.51 mg/kg has been reported (Shetty and Bhat 1997). (Da silva *et al.* 2004) reported that production of fumonisins by *Fusarium* species in Brazilian sorghum. Fumonisin have also been detected at lower levels in sorghum (Shetty and Bhat, 1997; Leslie and Marasas, 2001).

Consumption of cereals contaminated with fumonisins involves high health risks to animals and

humans. Recent studies also revealed that, fumonisins are associated with human oesophageal cancer. The International Agency for Research on Cancer (IARC) evaluated these toxins as human carcinogens (Fandohan *et al.* 2003). In view of their toxic properties, the FDA has fixed 2-4ppm/Kg of food as the permissible limit for fumonisins in foods. In view of all these, accurate detection and quantification of fumonisins in cereal samples is very important. It is also essential for a reliable evaluation of human exposure to these carcinogenic mycotoxins. Such detection systems set the realistic tolerance levels of fumonisins in food products for human and animal consumption. The data on the natural occurrence of fumonisins in cereals is also very much required for food and nutrition specialists for an appropriate toxicological evaluation. The study revealed the occurrence fumonisin producing strains of *Fusarium* species in many samples of maize and sorghum, warrants the need for systematic investigations of foods that are routinely channeled for human and animal consumption.

In conclusion, the research presented here demonstrates that TLC could reliably be used as an alternative to the more expensive methods of analyzing grain samples for fumonisins. Diagnostically, this TLC procedure can be used to initially screen sorghum and maize samples intended for human and animal consumption for the possible contamination with fumonisins, thus reducing the number of samples requiring quantitative analysis for the fumonisins by HPLC analysis in the laboratory. The present study is a sample investigation and further efforts are very much needed to know the epidemiological implications of these toxins on humans and animals. More studies are required to address this problem in other states of India.

REFERENCES

- Bailly, J.D.; Querin, A.; Tardieu, D. & Guerre, P. 2005. Production and purification of fumonisins from a highly toxigenic *Fusarium verticilloides* strain. *Revue de Medecine Veterinaire* 156(11): 547-554.
- Bhat, R.V.; Prathap Kumar, H.S.; Rao, P.A. & Rao, V.S. 1997. A food-borne disease outbreak due to the consumption of moldy sorghum and maize containing fumonisin mycotoxins. *Clinical Toxicology* 35:249-255. <http://dx.doi.org/10.3109/15563659709001208> PMID:9140318
- Chatterjee, D. & Mukherjee, S.K. 1994. Contamination of Indian maize with fumonisin B1 and its effects on chicken macrophage. *Letters in Applied Microbiology* 18:251-253. <http://dx.doi.org/10.1111/j.1472-765X.1994.tb00861.x>
- Chu, F.S. & Li, G.V. 1994. Simultaneous occurrence of fumonisin B₁ and other mycotoxins in moldy corn collected from People's Republic of China in regions with high incidence of esophageal cancer. *Applied and Environmental Microbiology* 60:847-852. PMID:8161178 PMCID:201401
- Da Silva, J.B.; Dilkin, P.; Fonseca, H. & Correa, B. 2004. Production of aflatoxin by *Aspergillus flavus* and fumonisin by *Fusarium* species isolated from Brazilian sorghum. *Brazilian Journal of Microbiology* 35:182-186.
- Fandohan, P.; Hell, K.; Marasas, W.F.O. & Wingfield, M.J. 2003. Infection of maize by *Fusarium* species and contamination with fumonisin in Africa. *African Journal of Biotechnology* 2:570-579.
- Gelderblom, W.C.A.; Seier, J.V.; Snijman, P.W.; Van Schalkwyk, D.J.; Shepard, G.S. & Marasas, W.F.O. 2001. Toxicity of culture material of *Fusarium verticillioides* strain MRC 826 to nonhuman primates. *Environment and Health Perspectives* 109:267-276. <http://dx.doi.org/10.1289/ehp.01109s2267> PMID:11359695 PMCID:1240675
- Ghianian, S.A.; Rezayat, S.M.; Kord-bacheh, P.; Maghsood, A.H.; Yazdanpanah, H.; Shephard, G.S.; Westhuizen, L.; Vismer, H.F. & Marasas, W.H.O. 2005. Fumonisin production by *Fusarium* species isolated from freshly harvested corn in Iran. *Mycopathologia* 159:31-40. <http://dx.doi.org/10.1007/s11046-004-3899-5> PMID:15750730
- Glenn, A.E. 2007. Mycotoxigenic *Fusarium* species in animal feed. *Animal Feed Science and Technology* 137(2007):213-240. <http://dx.doi.org/10.1016/j.anifeedsci.2007.06.003>
- Jindal, N.; Mahipal, S.K. & Rottinghaus, G.E. 1999. Occurrence of fumonisin B1 in maize and poultry feeds in Haryana, India. *Mycopathologia* 148:37-40. <http://dx.doi.org/10.1023/A:1007162113368> PMID:11086483
- Leslie, J.F. & Summerell, B.A. 2006. The *Fusarium*, Laboratory Manual. 1st ed. Blackwell Professional, USA. <http://dx.doi.org/10.1002/9780470278376>
- Leslie, J.F. & Marasas, W.F.O. 2001. *Fusarium* from sorghum: Life in interesting times. p.76-83. Proceeding of the 22nd biennial grain sorghum research and utilization conference, Nashville, Tennessee, USA.
- Logrieco, A.; Mule, G.; Moretti, A. & Bottalico, A. 2002. Toxigenic *Fusarium* species and mycotoxins associated with maize ear rot in Europe. *European Journal of Plant Pathology* 108:597-609. <http://dx.doi.org/10.1023/A:1020679029993>
- Morales-Rodriguez, I.; Yanez-Morales, M.J.; Silva-Rojas, H.V.; Garcia-de-los-Santos, G.; Guzman-de-Pena, D.A. 2007. Biodiversity of *Fusarium* species in Mexico associated with ear rot in maize, and their identification using a phylogenetic approach. *Mycopathologia* 163:31-39. <http://dx.doi.org/10.1007/s11046-006-0082-1> PMID:17216329 PMCID:2780687
- Munkvold, G.P.; Hellmich, R.L. & Showers, W.B. 1997. Reduced *Fusarium* ear rot and symptomless infection in kernels of maize genetically engineered for European corn borer resistance. *Phytopathology* 87:1071-1077. <http://dx.doi.org/10.1094/PHYTO.1997.87.10.1071> PMID:18945043
- Nakaya, S.C.; Shankar, A.C.U.; Niranjana, S.R.; Wulff, E.G.; Mortensen, C.N.; Prakash, H.S. 2010. Detection and quantification of fumonisins from *Fusarium verticillioides* in maize grown in southern India. *World Journal of Microbiology and Biotechnology* 26:71-78. <http://dx.doi.org/10.1007/s11274-009-0144-x>
- Nelson, P.E.; Plattner, R.D.; Shackelford, D.D.; Desjardins, A.E. 1992. Fumonisin B₁ production by *Fusarium* species other than *F. moniliforme* in Section *Liseola* and by some related species. *Applied and Environmental Microbiology* 58:984-989. PMID:1575501 PMCID:195366
- Rheeder, J.P.; Marasas, W.F.O. & Vismer, H.F. 2002. Production of fumonisin analogs by *Fusarium* species. *Applied and Environmental Microbiology* 68:2101-2105. <http://dx.doi.org/10.1128/AEM.68.5.2101-2105.2002> PMID:11976077 PMCID:127586
- Rheeder, J.P.; Marasas, W.F.O.; Thiel, P.G.; Sydenham, E.W.; Shephard, G.S. & Van schalkwyk, D.J. 1992. *Fusarium moniliforme* and fumonisins in corn in relation to human esophageal cancer in Transkei. *Phytopathology* 82:353-357. <http://dx.doi.org/10.1094/Phyto-82-353>
- Shaner, G. 2003. Epidemiology of *Fusarium* head blight of small grain cereals in North America. In: Leonard K, Bushnell WR, eds. *Fusarium* head blight of wheat and barley. St. Paul, Minnesota: APS Press. 84 p.
- Shephard, G.S.; Thiel, P.G.; Stockenstrom, S. & Sydenham, E.W. 1996. Worldwide survey of fumonisin contamination of corn and corn-based products. *Journal of Associative Official Analytical Chemistry International* 79:671-687.
- Shetty, P.H. & Bhat, R.V. 1997. Natural occurrence of fumonisin B1 and its co-occurrence with aflatoxin B1 in Indian sorghum, maize and poultry feeds. *Journal of Agriculture and Food Chemistry* 45:2170-2173. <http://dx.doi.org/10.1021/jf960607s>
- Sreenivasa, M.Y.; Dass, R.S.; Charith Raj, A.P. & Janardhana, G.R. 2006. Molecular detection of fumonisin producing *Fusarium*

species of freshly harvested maize kernels using Polymerase Chain Reaction (PCR). *Taiwania* 51: 1-10.

Sreenivasa, M.Y.; Dass, R.S.; Charith Raj, A.P. & Janardhana, G.R. 2008a. PCR method for the detection of genus *Fusarium* and fumonisin-producing isolates from freshly harvested sorghum grains grown in Karnataka, India. *Journal Food Safety* 28:236-247. <http://dx.doi.org/10.1111/j.1745-4565.2008.00117.x>

Sreenivasa, M.Y.; Gonzalez-jaen, M.T.; Dass, R.S.; Charith Raj, A.P. & Janardhana, G.R. 2008b. PCR-based assay for the detection and differentiation of potential fumonisin producing *F. verticillioides* isolated from Indian maize kernels. *Journal Food Biotechnology* 22(2):160-170. <http://dx.doi.org/10.1080/08905430802043206>

Tseng, T.C.; Tu, J.C. & Soo, L.C. 1995. Natural occurrence of mycotoxins in *Fusarium* infected beans. *Microbios* 84:21-28. PMID:8569524

Vismer, H.F.; Snijman, P.W.; Marasas, W.H.O. & Schalkwyk, D.J. 2004. Production of fumonisins by *Fusarium verticillioides* strains on solid and in a defined liquid medium - Effects of L-methionine and inoculum. *Mycopathologia* 158:99-106. <http://dx.doi.org/10.1023/B:MYCO.0000038428.06863.c2> PMID:15487327

Submit your next manuscript to IJLS with a -

- i) Convenient online submission,
- ii) Rapid editorial review followed by peer review,
- iii) No space constraints or color figure charges,
- iv) Immediate publication on acceptance.

Note1 - Authors/contributors are responsible for originality, contents, correct references, and ethical issues.

Note2 - Authors/contributors are required to write their text in good English (American or British usage but not a mixture of these), and authors who need assistance with language editing and copyediting services pre-/post-submission please visit <http://improveconsultant.com/> for more information.

© International Journal of Life Sciences (IJLS)